



Towards European Best Practice in Aquaculture Biofouling

A workshop organised by the EU-funded Collective Research on Aquaculture Biofouling and held during AQUA 2006, in Firenze, Italy

AQUA2006 – the combined meeting of the European Aquaculture Society and the World Aquaculture Society - brought together 3.250 participants from 86 countries in a scientific conference, a trade exhibition and several science-industry events. The European Commission, apart from having a pavilion in the exhibition, used AQUA2006 as a showcase for several EU-funded initiatives.

The CRAB project (Collective Research in Aquaculture Biofouling) is one of those initiatives focused on during AQUA2006. Apart from the workshop summarised in this document, members of the CRAB consortium also presented papers in the scientific conference and a CRAB stand in the exhibition provided a source of more detailed information to AQUA2006 participants. 34 participants from 19 countries attended the workshop. Analysis of organisation name showed that attendance was split 44% from industry (including industry associations), 44% from science and education and 12% from government institutions.

The full workshop presentations are available in PDF format on the CRAB web site (www.crabproject.com).

The workshop was split into two major sessions, preceded by a general introduction to biofouling and the CRAB initiative, and followed by a presentation of the dissemination and training aspects of the project. The first session addressed the impact of biofouling on infrastructure – notably net cages, platforms, ropes, buoys and all other equipment in contact with water. The second focused on fouling of the stock species, especially shellfish and the different set of problems that this poses. Each session was comprised of a presentation by one of the RTD partners in CRAB on the status of the research being carried out, followed by a presentation from one of the SME partners on the aspects that are important to industry. The floor was then open to participants for discussion on the CRAB approach and to compare the issues

in other countries.

The CRAB initiative, biofouling species in European aquaculture and associated problems

Peter Willemsen, Netherlands Organisation for Applied Scientific Research (TNO) & CRAB coordinator, The Netherlands.

Peter presented the general issues concerning biofouling and its effects on the European coastal aquaculture industry. This includes the general process, the biofouling organisms frequently encountered, the problem areas and estimations of the cost effects on the sector.

Biofouling is a complex and recurring problem in all sectors of the European aquaculture industry. Problem areas include biofouling on INFRASTRUCTURE (Immersed structures such as cages, netting and pontoons; equipment and structures such as pipelines, pumps, filters and holding tanks) and FARMED SPECIES (mussels, scallops, oysters etc). In the next 10 years the choice and availability of biocides for use as antifoulants will become much more restrictive within Europe with the application of the Biocides Products Directive EC 98/8/EC. Better knowledge within the sector will allow farmers to make informed choices and seek sustainable alternatives to current toxins that can cause pollution problems.

CRAB is reviewing current fouling control techniques and technologies and selecting (and optimising) suitable strategies to combat biofouling in aquaculture. These include biological control (using natural grazers); new materials such as non-toxic antifouling coatings; electrical methods (generating biocides (Cl-) or pH shifts) and new shellfish handling and immersion techniques.

Improving management practice and the possibility to predict



The effects of biofouling on infrastructure and on stock species are the two main pillars of the CRAB project.

biofouling will have advantages in the following production areas:

- Improving flow rates in cages and hence the growth rate of the fish
- Decreasing the frequency of net replacement and application of fouling products (coatings etc.)
- Reducing the stress on fish from net manipulation
- Reducing labour costs in shellfish cleaning

Especially this last point was raised for many SMEs in the shellfish sector. This cost can be the difference between profitability and failure for many small, often family-run businesses. But even for larger companies, the annual cost in net replacement and treatment for a medium sized salmon farm is estimated at €20k¹.

In presenting the CRAB initiative, Peter Willemsen pointed out that there is no “one size fits all” solution to biofouling. CRAB will contribute to the knowledge base by bringing together information that in the past has remained fragmented and rather inaccessible. He emphasized the importance of learning what may be useful from developments in other sectors (notably the shipping and pleasure boat sectors) for cost-effective application in aquaculture. He concluded by stating that more effective control and management of the biofouling problem will benefit the industry, with potential savings being 5-10% of the production value. The results from CRAB will be provided to the sector in the form of various tools that should be integrated in the overall production management protocols.

Reducing biofouling on infrastructure and equipment

Simone Dürr, University of Newcastle Upon Tyne, UK and **Donald Fowler**, Boris Net Company Ltd, UK

Simone and Donald co-presented a more detailed review of the CRAB strategies related to infrastructure, especially focussing on nets, and the possibilities offered by the use of slow-release anti-fouling coatings.

The three main biofouling principles are to combat settlement in the first place, to prevent fouling from developing and/or to remove the biofouling (by cleaning or reducing the adhesion forces).

Net cleaning, through various methods, is the most widely used strategy in many finfish producing countries, and may be done by the company itself, or, as in the case in Norway, by specifically trained companies contracted by producers.

The cleaning may be done underwater, or on land, after drying of the nets to kill the biofouling organisms.

Net cleaning is one thing, but net coatings are also important, especially new developments in non-toxic fouling release coatings. CRAB is testing some of these in “real-situation” farms.

However, the most important factor in managing biofouling remains the same in all locations. It is the possibility to accurately predict the occurrence of fouling episodes, especially mussel spat fall.

The discussion in this part of the workshop allowed the possibility for the CRAB partners present to exchange information with participants from European and non-European countries on the fouling problem and how to reduce fouling on infrastructure and equipment.

In Chile, fouling problems are mainly caused by tunicates, mussels and algae. The fouling problem is much more severe than in other countries due to the high productivity providing ideal bio-fouling conditions. In Chilean finfish farming, standard practice is to use Copper-oxide anti fouling treated nets, which are then removed from the cages and washed on-shore in industrial net washers. Recently, rigid and semi-rigid nettings for cages are being introduced into Chile; nettings such as ‘Aquagrid’ and metal mesh nettings, these do not currently require antifoulants. The main reason for their introduction is better predation deterrence. They are cleaned *in situ* as they are heavy and more difficult to handle, compared to conventional nylon netting. However, fish welfare may be improved due to the anti-predator properties of these nettings, possibly creating less stress. Furthermore, (salmon) escapes are a big issue in Chile (as in other salmon-



Various strategies are used for controlling biofouling on aquaculture nets



Non-toxic fouling release coatings may show promise for the finfish sector in Europe

¹ Own estimation by CRAB SME partners

producing countries, and nets that can prevent escapes are being tested. In Norway, the increase in cod production is accompanied by important efforts to reduce escapes, which cod seem particularly adept at doing. Cod eat their way through nylon mesh netting (as do sea bream in the Mediterranean), particularly at the base of nets. The use of polyester or 'dyneema' mesh can help eliminate this issue.

The preferred cleaning method for 'Aquagrid' is the underwater disk-cleaner. Cleaning has to be carried out on a regular 3-4 week cycle otherwise nets become badly fouled. Cleaning also provides improved welfare conditions. Labour costs for cleaning could be an issue in higher wage economies. In addition, these new nettings are much more expensive. Cost/benefit is therefore an important issue in appraising their efficiency.

In Canada (as in Chile) the major fouling organisms suggested as the main biofouling problem species in Canada are tunicates (*Ciona intestinalis*), algae, mussels and barnacles. The usual approach in fish farming is to use copper treated netting as antifoulant (as in Chile). The allowed copper incorporation rate is 20% higher than the levels used in European aquaculture. Air drying of nets is also used. Poly-culture techniques are currently being investigated. Research is showing that mussel culture introduced close to finfish farms may help to reduce settlement of tunicates.

From **Brazil** (from a shrimp farmer - this farm type is not included in the CRAB project). Participants indicated fouling in shrimp pools are mainly on the aeration paddle wheel, and this problem is solved by air drying and cleaning. In **Australia**-shrimp farming: participants indicated fouling on the paddle wheels was an issue primarily caused by barnacles and *Ulva* species.

In **New Zealand**, the fouling problems are caused by tunicates. The participant feels that invasive species will become more important in the near future.


In conclusion, the CRAB findings so far have been shown to be valid also for other countries. Similar solutions in finfish farming are being tried in Canada and Chile. Major fouling organisms seem to be tunicates, mussels and algae. Approaches against the problem are various and seem to be country/region and equipment specific.

Reducing biofouling in shellfish cultivation

Douglas Watson, University College Cork, Ireland and **John Icely**, Sagres, Portugal.

Douglas presented the status of CRAB work on shellfish biofouling. Antifouling measures are based on husbandry and cleaning methods and involve manual or mechanical cleaning, facilitated by dipping (for example in hot water) and air drying.

There is of course a crossover between the fouling on the shellfish themselves, and the fouling of infrastructure, buoys, lines, trays, lantern nets etc. mentioned in the previous section. Once again, the timing of any treatment is of primary importance.



(Photo courtesy of Douglas Watson, University College) Cork

However, CRAB is also assessing other strategies, including the use of grazers such as sea urchins (*Paracentrotus lividus*) and sea snails (*Monodonta lineata*) – shown above – in shellfish trays.

The possibility of allowing tunicates to foul shellfish is also being explored. The tunicates would prevent other fouling organisms taking hold, and are easy and therefore cheap to remove.

Furthermore, special shellfish coatings are being looked into for their efficacy.

John Icely, an oyster producer based on the Algarve coast in Portugal, summed up in simple words the costs and benefits of being involved in CRAB. He presented some basic data on the time (personnel cost) that is spent in cleaning oysters prior to being sold, or being re-stocked.

When management fails, one day's labour can easily become 6 times the effort. This translates into one person day at €35 becoming €210. For John and many other family-owned shellfish producers across Europe, the potential is high and may even represent actual economic viability in the sector.

The exchange between workshop participants from the shellfish sector gave rise to the following observations:

What happens when management fails?




Time to process nets increases
 Time to prepare shellfish for sale increases
 Time to clean lantern nets increases

1 days labour >>> 6 days labour
 One person day at 35 Euros >> 210 Euros

Biofouling is effectively a management issue

In **Australia** (one Pacific Oyster [*Crassostrea gigas*] farmer was present) the problem is not a particularly important one. Most of the farming of oyster there is carried out at intertidal sites and, due to the weather and the frequent turning of bags/infrastructure, the fouling on the shells and the material infrastructure does not appear to become pronounced. These conditions separate it from other parts of the world as they provide conditions, which are not ideal for extensive fouling. Any fouling on the equipment could easily be removed while the structures are being handled.

In addition to the intertidal nature of the farms contributing to reduced fouling, these sites also employ techniques such as cool rooms and immersion techniques to control fouling on the shellfish. Such techniques (particularly immersion techniques in warmer water – approx. 60°C) are also employed by CRAB SME partners, such as Sagres in Portugal.

In **Mexico** most shellfish cultivation occurs in the North East and North West of the country. No further details of the actual types of production or the problems facing the industry in this area were given by those who participated in the workshop. However, due to its proximity to the U.S.A. it is likely that production species and problems will be similar.

Sandra Shumway (an expert in bivalves and their biology, ecology, physiology and cultivation) was present at the meeting. She was able to give some insights into the financial issues facing the **North American** sector. She stated that fouling on the shellfish can be attributed to as much as 80% of the production costs within the industry. This is a huge figure, but is in fact likely to be comparable to other areas of the world, when the analysis is made. North America is primarily a site for the production of Pacific Oyster (*Crassostrea gigas*) and King Scallop (*Pecten maximus*).

European best practice guidelines and training activities

Alistair Lane, European Aquaculture Society

Alistair Lane presented the activities of the EAS and AquaTT in the dissemination and training tasks of CRAB. EAS is responsible for dissemination and will (with the other Association partners) build the Best Practice Guidelines based on findings of the work packages within CRAB, which will enable producers to select the most suitable strategy for their farming conditions.

AquaTT is responsible for building the training elements of CRAB. These include a biofouling manual and an online version of the Best Practice Guidelines. The combination of the two will allow a producer to consult regional information that will help in the prediction of biofouling events. The training tools will be developed during the Spring of 2007



CRAB – Collective Research on Aquaculture Biofouling – receives financial support from the Commission of the European communities, specific RTD programme “Collective Research” under contract COLL-CT-2003-500536. It does not necessarily reflect its views and in no way anticipates the commission’s future policy in this area”.



AQUA NOR Forum 2007

WELFARE-DRIVEN TECHNOLOGY

August 15-16, 2007

How our shared understanding of aquatic animals’ needs will shape their future homes.

A forum for consumers, industry, science and policy makers that discusses welfare as a driver for technological development in aquaculture.

Organised by the European Aquaculture Society, the Nor Fishing Foundation and SINTEF

More information at: www.easonline.org

and will be disseminated throughout European aquaculture operations soon after. The will of course be posted on the CRAB web site (www.crabproject.com).

Concluding comments

While attendance was limited in numbers, it was certainly not in terms of the number of countries represented. This gave a unique opportunity to exchange information with other regions and countries and to provide feedback on the success of various strategies.

The workshop clearly indicated a selection of strategies clearly linked to national legislation. In the European context, the EU Directives are of course translated into Member State legislation. Different countries therefore have the possibility to select antifouling strategies (and compounds) that other countries do not have access to.

The workshop also gave the possibility for the Collective Research tool itself to be presented. There is a need (and a desire) for European RTD to move beyond Europe, but at the same time, Collective Research is targeted towards improving European competitiveness. The structure of CRAB, and the challenges that the CRAB consortium faces in terms of the success of the initiative, show that European initiatives such as CRAB are useful as a model for cooperation in other countries and regions.

